arXiv:0912.1287v1 [quant-ph] 7 Dec 2009

Nonadiabatic tunneling in an ideal one dimensional semi-infinite periodic potential systems: An analytically solvable model

Aniruddha Chakraborty
Department of Inorganic and Physical Chemistry,
Indian Institute of Science, Bangalore, 560012, India.

(Dated: December 7, 2009)

Nonadiabatic tunneling in an ideal one dimensional semi-infinite periodic potential system is analyzed using our model. Using our simple analytically solvable model it is shown that ideal one dimensional semi-infinite periodic potential system can be thought as a model of moleular switch. The method is applicable to those systems where Green's function for the motion in the absence of coupling is known. The method has been applied to the problem where motion takes place on parabolic potentials, for which analytical expression of Green's function can be found.

INTRODUCTION

Quantum mechanical tunneling is an extremely important phenomenon in various fields of physics, chemistry and biology [1]. Most of the electronic transitions in molecular systems can be understood as non adiabatic transitions among electronically adiabatic eigen states at each fixed nuclear configuration. Nonadiabatic tunneling, is a tunneling accompanied by a nonadiabatic transition [2, 3, 4]. The nonadiabatic tunneling is different from ordinary tunneling which involves single potential. It should be noted that these two cases become the same under strong diabatic coupling limit. This is not the case in general, and the tunneling is affected by the upper adiabatic potential. So the nonadiabatic tunneling process can be controlled by the diabatic coupling, i.e., this can be switched on and off by switching the coupling. In this paper we are dealing with nonadiabatic tunneling in a one dimensional semi-infinite periodic potential system. Switching in one dimensional periodic potential systems have been gaining attention since the original works done by Kronig and Penney [5]. Complete transmission, which occurs at certain discrete energies in a semi-infinite periodic potential energy system, can be completely switched off by using a complete reflection due to an impurity potential created in the array. Positions of the complete reflections are determined by the shape of the potential and the nature of coupling. So this complete switching can be realized by changing one of the potential in the array to create an impurity. Similar idea was proposed by Carter [6], based on an array of periodic potential barriers. The molecular switching model based on finite nonadiabatic tunneling potential system was originally proposed by Nanbu *et al.* [7]. In thise paper we use one dimensional semi-infinite periodic potential systems to show the possibility of molecular switching using our simple analytically solvable model. In the following we will first discuss our two state model.

OUR TWO STATE MODEL

We consider two diabatic curves, crossing each other. There is a coupling between the two curves, which causes transitions from one curve to another. This transition would occur, in the simplest approximation near the vicinity of the crossing point. In particular, it will occur in a narrow range of x, given by

$$V_1(x) - V_2(x) \simeq V_{12}(x_c) \tag{1}$$

where x denotes the nuclear coordinate and x_c is the crossing point. V_1 and V_2 are determined by the shape of the diabatic curves and V_{12} represent the coupling between them. Therefore it is interesting to analyse a model, where coupling is localized in space near x_c. Thus we put

$$V_{12}(x) = K_0\delta(x - x_c) \tag{2}$$

here K_0 is a constant. This model has the advantage that it can be exactly solved [8, 9, 10]. In the following we will first discuss our result of the two state curve crossing problem, which will be used in later section to analyze an ideal one dimensional semi-infinite periodic potential system.

TWO STATE CURVE CROSSING PROBLEM: EXACT ANALYTICAL RESULTS FOR THE DELTA FUNCTION COUPLING

We start with a particle moving on any of the two diabatic curves and the problem is to calculate the probability that the particle will still be in that diabatic curve after a time t. We write the probability amplitude for the particler as

$$\Psi(x,t) = \begin{pmatrix} \psi_1(x,t) \\ \psi_2(x,t) \end{pmatrix} \tag{3}$$

Where $\psi_1(x,t)$ and $\psi_2(x,t)$ are the probability amplitude for the two states. $\Psi(x,t)$ obey the time dependent Schrӧdinger equation (we put $\hbar = 1$)

$$i\frac{\partial \Psi(x,t)}{\partial t} = H\Psi(x,t) \tag{4}$$

In the above equation H is defined by

$$H = \begin{pmatrix} H_1(x) & V_{12}(x) \\ V_{21}(x) & H_2(x) \end{pmatrix} \tag{5}$$

where $H_1(x)$, $H_2(x)$ and $V_{12}(x)$ are

$$H_1(x) = -\frac{1}{2m}\frac{\partial^2}{\partial x^2} + V_1(x) \tag{6}$$

$$H_2(x) = -\frac{1}{2m}\frac{\partial^2}{\partial x^2} + V_2(x) \tag{7}$$

$$V_{12}(x) = V_{21}(x) = V(x) \tag{8}$$

In the above $V_1(x)$ and $V_2(x)$ are determined by the shape of that diabatic curve. $V(x)$ is a position dependent coupling function we shall eventually assume that $V(x)$ is a Dirac delta function. In the following we will use the abstract operator notation of quantum mechanics. We found it convenient to define the half Fourier Transform $\overline{\Psi}(E)$ of $\Psi(t)$

$$\overline{\Psi}(E) = \int_0^\infty \Psi(t) e^{iEt} dt \tag{9}$$

Half Fourier transformation of Eq. (4) gives

$$\begin{pmatrix} \overline{\psi}_1(E) \\ \overline{\psi}_2(E) \end{pmatrix} = i \begin{pmatrix} E - H_1 & V \\ V & E - H_2 \end{pmatrix}^{-1} \begin{pmatrix} \psi_1(0) \\ \psi_2(0) \end{pmatrix} \tag{10}$$

This may be written as

$$\overline{\Psi}(E) = iG(E)\Psi(0) \tag{11}$$

$G(E)$ is defined by $(E-H)\ G(E) = I$. In the position representation, the above equation may be written as

$$\overline{\Psi}(x,E) = i\int_{-\infty}^{\infty} G(x,x_0;E)\overline{\Psi}(x_0,E)dx_0 \tag{12}$$

The Green's function is defined by

$$G(x,x_0;E) = \langle x|(E-H)^{-1}|x_0\rangle \tag{13}$$

where $|x_0\rangle$ denotes the position eigen ket with an eigenvalue x_0

$$G(x,x_0;E) = \begin{pmatrix} G_{11}(x,x_0;E) & G_{12}(x,x_0;E) \\ G_{21}(x,x_0;E) & G_{22}(x,x_0;E) \end{pmatrix} \tag{14}$$

Using the partition technique [11]we can write $G_{11}(x, x_0; E)$ as given below

$$G_{11}(x, x_0; E) = \langle x|[E - H_1 - V(E - H_2)^{-1}V]^{-1}|x_0\rangle \tag{15}$$

The above equation is true for any general V. This equation considerably simplifies if V is a Dirac delta function at x_c, which in the operator notation may be written as $V = K_0 S = K_0\ |x_c\rangle\ \langle x_c|$. Then

$$G_{11}(x, x_0; E) = \langle x|[E - H_1 - K_0^2 G_2^0(x_c, x_c; E)S]^{-1}|x_0\rangle \tag{16}$$

$G_2^0(x, x_0; E)$ is defined by

$$G_2^0(x, x_0; E) = \langle x|(E - H_2)^{-1}|x_0\rangle \tag{17}$$

and corresponds to the propagation of the particle starting from x_0, on the second surface in the absence of coupling. Now we will use the operator identity

$$(E - H_1 - K_0^2 G_2^0(x_c, x_c; E)S)^{-1} =$$

$$(E - H_1)^{-1} + (E - H_1)^{-1} K_0^2 G_2^0(x_c, x_c; E)S[E - H_1 - K_0^2 G_2^0(x_c, x_c; E)S]^{-1} \tag{18}$$

Inserting the resolution of identity $I = \int_{-\infty}^{\infty} dy\ |y\rangle\ \langle y|$ in the second term of the above equation, we arrive at a Lippman-Schwinger type equation

$$G_{11}(x, x_0; E) = G_1^0(x, x_0; E) + K_0^2 G_2^0(x_c, x_c; E) \int_{-\infty}^{\infty} dy G_1^0(x, y; E)\delta(y - x_c)G_{11}(y, x_0; E) \tag{19}$$

$$G_{11}(x, x_0; E) = G_1^0(x, x_0; E) + K_0^2 G_1^0(x, x_c; E)G_2^0(x_c, x_c; E)G_{11}(x_c, x_0; E) \tag{20}$$

we now put $x = x_c$ in Eq. (20) and solve it to find

$$G_{11}(x_c, x_0; E) = G_1^0(x_c, x_0; E)[1 - K_0^2 G_1^0(x_c, x_c; E)G_2^0(x_c, x_c; E)]^{-1} \tag{21}$$

This when substituted back into Eq. (20) gives

$$G_{11}(x, x_0; E) = G_1^0(x, x_0; E) + \frac{K_0^2 G_1^0(x, x_c; E)G_2^0(x_c, x_c; E)G_1^0(x_c, x_0; E)}{1 - K_0^2 G_1^0(x_c, x_c; E)G_2^0(x_c, x_c; E)} \tag{22}$$

By using the same procedure one can derive analytical expression of $G_{12}(x, x_0; E)$.

$$G_{12}(x, x_0; E) = \frac{K_0 G_1^0(x, x_c; E)G_2^0(x_c, x_0; E)}{1 - K_0^2 G_1^0(x_c, x_c; E)G_2^0(x_c, x_c; E)} \tag{23}$$

Similarly one can derive the expression for $G_{22}(x, x_0; E)$ and $G_{21}(x, x_0; E)$. Once $G(x, x_0, E)$ is known, one can calculate wave function explicitely and using the wave function one can easily calculate transition probability from one diabatic potential to the other. In the following, we will apply our two state result to the case of one dimensional semi-infinite periodic potential system.

ONE DIMENSIONAL SEMI-INFINITE PERIODIC POTENTIAL SYSTEM

We imagine the problem of atom hopping reaction on a particular surface. Suppose active sites (which can adsorb atom) on the surface are arranged on a straight line. So we are thinking of one dimensional motion of an atom along the surface. The potential energy curve (after fixing the position of active sites) as a function of the position of atom will look like "one dimensional periodic potential system". Now we consider the motion of active sites too. We take two neighboring active sites A and B, where an atom from A is transferred to B. The potential energy curve for $\overset{*}{A}\ B$ and $A\ \overset{*}{B}$ crosses at a point (see Fig. 1). When this same atom transfers from B to C (the next neighboring active site), the potential energy curve for $\overset{*}{B}\ C$ and $B\ \overset{*}{C}$ crosses at another point. Here we assume that the motion of all active sites are coupled to the motion of it's nearest neighbouring active sites via Dirac delta functions - so that we can use our two state result to analyze this problem. So if an atom transports from one active site to another active sites through n active sites on the surface, there will be $n + 1$ potential energy curves and n crossing points (see Fig. 2). Here we assume n to be very large (semi-infinite limit). In the following we will analyse this semi-infinite periodic potential system using our two state result.



FIG. 1: The potential energy curve for $\overset{*}{A}\,B$ (left) and $A\,\overset{*}{B}$ (right) crosses at a point.

GREEN'S FUNCTION FOR ONE DIMENSIONAL SEMI-INFINITE PERIODIC POTENTIAL SYSTEMS

Now first we assume that the all active sites are equivalent and the distance between any two neighboring active sites is the same, so the distance between any two neighboring crossing points will be the same. We have $(n+1)$ potential energy curves, we want to start our calculation with the n-th potential energy curve and let $G_0^n(x, x_0; E)$ denote the Green's function for the vibrational motion of the n-th active site, when it's vibration is uncoupled from the vibrations of all other active sites. Let $G^{(n-1)}(x, x_0; E)$ denote the Green's function for the vibrational motion of first $(n-1)$ active sites, when only the vibrational motion of $(n-1)$-th active site is uncoupled from that of the n-th active site. Now using our two state theory (coupling is represented by Dirac delta function) we have derived an analytical expression of the Green's function for the vibrational motion of first n active sites, where the potential energy curve describing the vibrational motion of n-th active site is coupled to that of $(n-1)$-th active site at x_{n-1}.

$$G^{(n)}(x, x_0; E) = G_0^n(x, x_0; E) + \frac{K_0^2 G_0^n(x, x_{n-1}; E) G^{(n-1)}(x_{n-1}, x_{n-1}; E) G_0^n(x_{n-1}, x_0; E)}{1 - K_0^2 G_0^n(x_{n-1}, x_{n-1}; E) G^{(n-1)}(x_{n-1}, x_{n-1}; E)} \tag{24}$$

Now we calculate the Green's function for the vibrational motion of the $(n+1)$ active sites, where the potential energy curve describing the vibrational motion of the $(n+1)$-th active site is coupled to that of the n-th active site at x_n. If n is very large (semi-infinite) using the symmetry argument one can easily prove that

$$G^{(n)}(x_n, x_n; E) = G^{(n-1)}(x_{n-1}, x_{n-1}; E) \tag{25}$$

So we get

$$G^{(n)}(x, x_0; E) = G_0^n(x, x_0; E) + \frac{K_0^2 G_0^n(x, x_{n-1}; E) G^{(n)}(x_n, x_n; E) G_0^n(x_{n-1}, x_0; E)}{1 - K_0^2 G_0^{n+1}(x_n, x_n; E) G^{(n)}(x_n, x_n; E)}. \tag{26}$$



FIG. 2: The potential energy curves as a function of vibrational motion of all active sites.

If we put $x = x_n$ and $x_0 = x_n$, in the above equation we get

$$G^{(n)}(x_n, x_n; E) = G_0^n(x_n, x_n; E) + \frac{K_0^2 G_0^n(x_n, x_{n-1}; E) G^{(n)}(x_n, x_n; E) G_0^n(x_{n-1}, x_n; E)}{1 - K_0^2 G_0^{n+1}(x_n, x_n; E) G^{(n)}(x_n, x_n; E)} \tag{27}$$

The above equation can be simplified to

$$\begin{gathered} -[K_0^2 G_0^{n+1}(x_n, x_n; E) G_0^n(x_n, x_n; E) - K_0^2 G_0^n(x_n, x_{n-1}; E) G_0^n(x_{n-1}, x_n; E) + 1] G^{(n)}(x_n, x_n; E) \\ K_0^2 G_0^{n+1}(x_n, x_n; E) G^{(n)}(x_n, x_n; E)^2 + G_0^n(x_n, x_n; E) = 0 \end{gathered} \tag{28}$$

Now we solve the above equation for $G^{(n)}(x_n, x_n; E)$.

$$\begin{aligned} G^{(n)}(x_n, x_n; E) = {} & \frac{[K_0^2 G_0^{n+1}(x_n,x_n;E) G_0^n(x_n,x_n;E) - K_0^2 G_0^n(x_n,x_{n-1};E) G_0^n(x_{n-1},x_n;E) + 1]}{2K_0^2 G_0^{n+1}(x_n,x_n;E)} \\ & \pm \frac{\sqrt{[K_0^2 G_0^{n+1}(x_n,x_n;E) G_0^n(x_n,x_n;E) - K_0^2 G_0^n(x_n,x_{n-1};E) G_0^n(x_{n-1},x_n;E) + 1]^2 - 4K_0^2 G_0^{n+1}(x_n,x_n;E) G_0^n(x_n,x_n;E)}}{2K_0^2 G_0^{n+1}(x_n,x_n;E)} \end{aligned} \tag{29}$$

In the above expression, $G^{(n)}(x_n, x_n; E)$ is expressed in terms of Green's functions for uncoupled potential. For a continuum of vibrational eigen energies $G^{(n)}(x_n, x_n; E)$ will have non-zero imaginary part for a range of E and that is possible only if the quantity $A(E)$ as defined below, is negative.

$$\begin{aligned} A = -4K_0^2 G_0^{n+1}(x_n, x_n; E) G_0^n(x_n, x_n; E) + [& K_0^2 G_0^{n+1}(x_n, x_n; E) G_0^n(x_n, x_n; E) \\ & - K_0^2 G_0^n(x_n, x_{n-1}; E) G_0^n(x_{n-1}, x_n; E) + 1]^2 \end{aligned} \tag{30}$$

So those enrgy ranges for which $A(E)$ is negative, complete transmission is possible. In the following we will discuss the result of our numerical calculation using Eq. (30).



FIG. 3: The plot of "A" against E near the the energy of zoro-th vibrational state.

APPLICATION TO THE HARMONIC POTENTIAL CASE

It is possible to find $G_0^n(x, x_0; E)$ only for few cases and harmonic potential is one of them. For example, if

$$V_1(x) = \frac{1}{2} m \omega_0^2 x^2 \tag{31}$$

then the analytical expression of the corresponding Green's function is known[12]. We calculate A for harmonic potentials [Eq. (30)]. We use harmonic potential of frequency $\omega_0 = 500\ cm^{-1}$ and mass $m = 35.4$. The distance between the equilibrium positions of any two neighboring harmonic potential we take, is $0.1\mathring{A}$. The distance between the crossing points of any two neighboring harmonic potential we take, is $0.05\mathring{A}$. K_0 we take is $1.58 \times 10^{-7}\ erg.\mathring{A}$ [8]. The result of our calculation (see Fig. 3) shows that "A" is negative over a range of energy near zero point vibration ($\frac{\omega_0}{2}$), this means that $G^{(n)}(x_n, x_n; E)$ has non-zero imaginary part over the same range of energy, so the system has a continuum of vibrational eigen energies in that range of energy. Same is true for other vibrations ($\frac{3\omega_0}{2}$, $\frac{5\omega_0}{2}$.......) also. So complete transmission is possible near those energy values ($\frac{\omega_0}{2}$, $\frac{3\omega_0}{2}$, $\frac{5\omega_0}{2}$.......). These complete transmissions can be completely switched off by using a complete reflection due to an impurity potential created in the array.

CONCLUSIONS:

Nonadiabatic tunneling in an ideal one dimensional semi-infinite periodic potential system is analyzed using our model Hamiltonian. It was shown that complete transmission is possible for certain range of energies and complete reflection is possible by adding impurity potential in the array. The same model can be applied to multi-state electron transfer model, where the atomic coordinate will be replaced by a global coordinate representing the nuclear polarization of the system. Our method is applicable to those systems, where Green's function for the motion in the absence of coupling is known. The method has been applied to the problem where motion takes place on parabolic potentials, for which the Green's function can be found. The same procedure is also applicable to the case where S is a non-local operator and may be represented by $S \equiv |f\rangle\ K_0\ \langle g|$, f and g are arbitrary acceptable functions. Choosing both of them to be Gaussian should be an improvement over the delta function coupling model. S may even be a linear combination of such operators.

ACKNOWLEDGMENTS:

The author thanks Prof. K. L. Sebastian for suggesting this very interesting problem. It is a pleasure to thank Prof. M. S. Child for his kind interest, suggestions and encouragement.

[1] R. P. Bell, *The tunneling effect in chemistry* (Chapman and Hall, London, 1980).
[2] H. Nakamura, *Int. Rev. Phys. Chem.*, **10**, *123* (1991).
[3] H. Nakamura, in *Theory, Advances in Chemical Physics*, edited by M. Bayer and C. Y. Ng (John Wiley and Sons, New York, 1992).
[4] H. Nakamura, *Nonadiabatic Transition*, (World Scientific, Singapore, 2002).
[5] R. de L. Kronig and W. G. Penney, Proc. R. Soc. (London), Ser. A **130**,*499*(1931).
[6] F. L. Carter *Molecular Electronic Devices*,(Marcel-Dekker, New York, 1982).
[7] S. Nanbu, H. Nakamura and F. O. Goodman, J. Chem. Phys. **107**, *5445* (1997).
[8] A. Chakraborty, Mol. Phys. **107**, *165* (2009).
[9] A. Chakraborty, Mol. Phys. (in press).
[10] A. Chakraborty, Mol. Phys. (under revision).
[11] P. Lowdin, J. Math. Phys. **3**, *969* (1962).
[12] C. Grosche and F. Steiner, *Handbook of Feynman Path Integral* (Springer-Verlag, Berlin, 1998).